

ON DOUBLE COVERINGS BETWEEN SMOOTH PLANE CURVES

TAKESHI HARUI, JIRYO KOMEDA AND AKIRA OHBUCHI

ABSTRACT. We completely classify the pairs of two smooth plane curves with double coverings between them. More precisely, we show that there exist no double coverings between two smooth plane curves except for several special cases.

1. INTRODUCTION AND PRELIMINARIES

We consider the following problem on smooth plane curves:

Problem. Classify the pairs of smooth plane curves (C, C') such that there exists a covering $\pi : C \rightarrow C'$ of degree $n \geq 2$.

In this article we completely classify such pairs in the case $n = 2$. There exist several trivial pairs: $(\mathbb{P}^1, \mathbb{P}^1)$, (E, \mathbb{P}^1) and (E, E') , where E and E' are elliptic curves with an isogeny $\pi : E \rightarrow E'$ of degree 2. We give another example.

Example 1.1. Let C be a Fermat quartic defined by $x^4 + y^4 = 1$. It has an involution $\sigma : C \rightarrow C((x, y) \mapsto (-x, y))$. Then $B = C^{(\sigma)}$ is an irreducible curve defined by $x^2 + y^4 = 1$ and $\tau : C \rightarrow B((x, y) \mapsto (x^2, y))$ is a double covering. Note that the normalization E of B is an elliptic curve and there exists a lifted double covering $\pi : C \rightarrow E$. Thus we obtain a pair (C, E) of smooth plane curves with a double covering from C to E .

In fact, we show that pairs of the above type (a smooth plane quartic with an involution and an elliptic curve) are the only non-trivial ones (see Theorem 2.1 and Proposition 2.6).

Notation and Conventions

A curve is a one-dimensional algebraic variety. It is possibly reducible or non-reduced, though it will be smooth and irreducible mainly in this article. A g_d^r is a linear system of degree d and dimension r on a smooth curve. It is said to be *simple* if the associated rational map to \mathbb{P}^r is birational onto its image. Otherwise it is said to be *compounded*.

For two divisors D and D' on a variety, $D \sim D'$ denotes that they are linearly equivalent.

2. RESULTS

Our main result is the following theorem.

Theorem 2.1. *Let C, C' be two smooth plane curves of degree d, d' , respectively. Then there exist no double coverings from C to C' except for the following cases:*

- (1) C' is rational ($d' \leq 2$) and C is rational or elliptic ($d \leq 3$).
- (2) C and C' are elliptic ($d = d' = 3$).
- (3) C' is elliptic and C is a bi-elliptic plane quartic ($d' = 3, d = 4$).

In particular, a double covering of a smooth plane curve of degree greater than three is never a smooth plane curve.

Our proof of this theorem consists of two parts. First we consider the case of small degree.

Lemma 2.2. *Let C, C' be two smooth plane curves of d, d' , respectively. Assume that $d' \leq 3$ and there exists a double covering $\pi : C \rightarrow C'$. Then one of the following holds:*

- (1) C' is rational ($d' \leq 2$) and C is rational or elliptic ($d \leq 3$).
- (2) C and C' are elliptic ($d = d' = 3$).
- (3) C' is elliptic and C is a bi-elliptic plane quartic ($d' = 3, d = 4$).

Proof. First assume that C' is rational. Then C is hyperelliptic. On the other hand, it is well-known that $\text{gon}(C) = d - 1$. Hence we have $d \leq 3$, which implies that C is rational or elliptic. Secondly, assume that C' is elliptic. Then C is bi-elliptic. It is known that $s(2) \geq g + 1$ for a bi-elliptic curve of genus g . On the other hand, $s(C, 2) = d$ holds because C is a smooth plane curve of degree d . Thus we have

$$d = s(C, 2) \geq g(C) + 1 = \frac{1}{2}(d - 1)(d - 2) + 1,$$

which implies that $(d - 1)(d - 4) \leq 0$. Hence $d \leq 4$ holds. \square

Next we show the non-existence in the case of large degree.

Proposition 2.3. *Let C, C' be two smooth plane curves of d, d' , respectively. If $d' \geq 4$, then there exist no double coverings from C to C' .*

Proof. Assume that $d' \geq 4$ and there exists a double covering $\pi : C \rightarrow C'$. First note that C' has two pencils of degree $d' - 1$ and d' . The pull-back of them by π are pencils of degree $2(d' - 1)$ and $2d'$ on C . Since $\text{gon}(C) = d - 1$, we have $d - 1 \leq 2(d' - 1)$, i.e., $2d' \geq d + 1$. On the other hand, C has no pencils of degree between $d + 1$ and $2d - 5$ (cf. [C]). Hence we obtain that $2d' \geq 2d - 4$, i.e., $d' \geq d - 2$. Since $d > d'$ by our assumption $d' \geq 4$, we have $d' = d - 1$ or $d' = d - 2$.

Case(i) $d' = d - 1$. We then have

$$\begin{aligned} g(C) &= \frac{1}{2}(d-1)(d-2) \quad \text{and} \\ g(C') &= \frac{1}{2}(d'-1)(d'-2) = \frac{1}{2}(d-2)(d-3). \end{aligned}$$

Since $g(C) \geq 2g(C') - 1$ holds by Riemann-Hurwitz formula, we obtain that

$$\frac{1}{2}(d-1)(d-2) \geq (d-2)(d-3) - 1,$$

which implies that $(d-2)(d-5) \leq 2$. Hence we have $d \leq 5$. Thus it suffices to exclude the case $(d, d') = (5, 4)$. In this case there exists a divisor N on C' of degree 1 such that $C = \text{Spec}(\mathcal{O}_{C'} \oplus \mathcal{O}_{C'}(-N))$. Since $g(C') = 3$, the morphism $C'^3 \rightarrow \text{Pic}^1(C')((P, Q, R) \mapsto P + Q - R)$ is surjective. Hence $N \sim P + Q - R$ for some $P, Q, R \in C'$. We take an effective divisor D on C' linearly equivalent to $K_{C'} - R$. Then we have

$$\begin{aligned} h^0(C, \mathcal{O}_C(\pi^*D)) &= h^0(C', \mathcal{O}_{C'}(D)) + h^0(C', \mathcal{O}_{C'}(D - N)) \\ &= h^0(C', \mathcal{O}_{C'}(D)) + h^0(C', \mathcal{O}_{C'}(K_{C'} - P - Q)) \\ &= 2 + 1 = 3. \end{aligned}$$

Hence we obtain a net $|\pi^*D| = g_6^2$ on C . If this is compounded, then C is a bi-elliptic plane quintic, which leads to a contradiction by Lemma 2.2. If the g_6^2 is simple, then C has at most five g_4^1 's. This is again a contradiction, since a smooth plane quintic C has infinitely many g_4^1 's. Thus we complete the proof in this case.

Case(ii) $d' = d - 2$. In this case C has a pencil of degree $2(d' - 1) = 2d - 6$. Recall that C' has no pencils of degree between $d+1$ and $2d-5$. Hence we have $2d-6 \leq d$, which implies that $d \leq 6$. Thus we only have to consider the case $(d, d') = (6, 4)$. In this case we take a ramification point P of $\pi : C \rightarrow C'$ and set $P' := \pi(P)$. Then one of the following holds for the Weierstrass semigroup $H(P')$ of C' at P' :

$$H(P') = \begin{cases} \langle 3, 4 \rangle, \\ \langle 3, 5, 7 \rangle \quad \text{or} \\ \langle 4, 5, 6, 7 \rangle. \end{cases}$$

In the first two cases $H(P')$ contains 3 and 7, which implies that $H(P)$, the Weierstrass semigroup of C at P , contains 6 and 14. In the third case $H(P)$ contains 8, 10, 12 and 14. Then we obtain a contradiction in each case by using the lemma below. Thus we complete the proof. \square

Lemma 2.4. ([K], [KK]) *Let C be a smooth plane sextic.*

- (1) *If $H(P)$ contains 6 for some point $P \in C$, then $H(P) = \langle 5, 6 \rangle$.*
- (2) *There exists no point $P \in C$ such that $H(P)$ contains 8, 10, 12 and 14.*

Proof. For two plane curves Γ_1 and Γ_2 such that no common component of them contains P , we denote by $I(\Gamma_1 \cap \Gamma_2, P)$ the intersection multiplicity of Γ_1 and Γ_2 at P . Since plane cubics cut out the canonical linear system on C , we have

$$G(P) := \mathbb{N}_0 \setminus H(P) = \{I(C \cap \Gamma, P) + 1 \mid \Gamma \text{ is a plane cubic}\}.$$

Note that $\sharp G(P) = 10$. Let T be the tangent line to C at P , L_1 a line in \mathbb{P}^2 that is distinct with T and contains P and L_0 a line in \mathbb{P}^2 not containing P . For any non-negative integers α and β with $\alpha + \beta \leq 3$, we denote by $\Gamma(\alpha, \beta)$ the plane cubic $\alpha T + \beta L_1 + (3 - \alpha - \beta)L_0$. We can write $T|_C = \nu P + D$, where ν is an integer such that $2 \leq \nu \leq 6$ and D is an effective divisor on C whose support does not contain P .

(1) First note the following.

$$\text{If } \nu = 5, \text{ then we have } I(C \cap \Gamma(1, 0), P) = 5.$$

$$\text{If } \nu = 4, \text{ then we have } I(C \cap \Gamma(1, 1), P) = 5.$$

$$\text{If } \nu = 3, \text{ then we have } I(C \cap \Gamma(1, 2), P) = 5.$$

$$\text{If } \nu = 2, \text{ then we have } I(C \cap \Gamma(2, 1), P) = 5.$$

Thus we obtain that $G(P)$ contains 6 if $\nu \leq 5$. In other words, if $H(P)$ contains 6, then $\nu = 6$ holds. Then we have $I(C \cap \Gamma(\alpha, \beta), P) = 6\alpha + \beta$ for $0 \leq \alpha + \beta \leq 3$. It follows that

$$G(P) \supset \{1, 2, 3, 4, 7, 8, 9, 13, 14, 19\}.$$

Hence we obtain that $G(P) = \{1, 2, 3, 4, 7, 8, 9, 13, 14, 19\}$ by comparing the cardinality of these sets. Equivalently, $H(P) = \langle 5, 6 \rangle$ holds.

(2) First we exclude the case where $\nu \geq 3$.

$$\text{If } \nu = 6, \text{ then we have } I(C \cap \Gamma(1, 1), P) = 7.$$

$$\text{If } \nu = 5, \text{ then we have } I(C \cap \Gamma(1, 2), P) = 7.$$

$$\text{If } \nu = 4, \text{ then we have } I(C \cap \Gamma(2, 1), P) = 9.$$

$$\text{If } \nu = 3, \text{ then we have } I(C \cap \Gamma(2, 1), P) = 7.$$

Thus we obtain that $H(P) \not\cong 8$ or $H(P) \not\cong 10$ in these cases. Suppose that $\nu = 2$. Then there exists a conic B such that $i_2 := I(C \cap B, P) \geq 5$. Note that $i_2 \leq 12$. If $i_2 = 5$, then $I(C \cap (T + B), P) = 7$, which implies that $H(P) \not\cong 8$. If i_2 is even with $i_2 \leq 12$, then $I(C \cap (L_1 + B), P) = i_2 + 1$, which implies that $H(P) \not\cong i_2 + 2$. If i_2 is odd with $7 \leq i_2 \leq 11$, then $I(C \cap (L_0 + B), P) = i_2$, which implies that $H(P) \not\cong i_2 + 1$. Thus we obtain a contradiction in any case.

Remark 2.5. Weierstrass semigroups of smooth plane curves of low degree are classified (cf. [K], [KK]). Our argument in the proof of the above lemma is based on theirs.

□

In the end, we remark a fact on smooth plane quartics.

Proposition 2.6. *Let C be a smooth plane quartic. Then it is bi-elliptic if and only if it has an involution.*

Proof. It is clear that a bi-elliptic curve has an involution. Conversely, assume that C has an involution σ . We then have a double covering $\pi : C \rightarrow C'$, where C' is the normalization of the (possibly singular) curve $C^{(\sigma)}$. Then $g(C') = 0, 1$ or 2 . It is known that a curve of genus 3 with a double covering onto a curve of genus 2 is hyperelliptic (cf. [A]). Since C is non-hyperelliptic, we have $g(C') = 1$. Thus C is bi-elliptic. \square

REFERENCES

- [A] R. D. M. Accola: On lifting the hyperelliptic involution, Proc. Amer. Math. Soc. **122** (1994), 341-347.
- [ACGH] E. Arbarello, M. Cornalba, P. A. Griffiths, J. Harris: Geometry of Algebraic Curves Vol. I, Grundlehren 267, Springer-Verlag, 1985.
- [C] M. Coppens: The existence of base point free linear systems on smooth plane curves, J. Alg. Geom. **4** (1995), 1-15.
- [CK] M. Coppens, T. Kato: The gonality of smooth curves with plane models, Manuscripta Math. **70** (1990), 5-25.
- [CKM] M. Coppens, C. Keem, G. Martens: Primitive linear series on curves, Manuscripta Math. **77** (1992), 237-264.
- [HKO] T. Harui, K. Kato, A. Ohbuchi: The minimal degree of plane models of algebraic curves and double coverings, preprint.
- [HO] T. Harui, A. Ohbuchi: On the number of pencils of minimal degree on curves with small gonality, preprint.
- [K] S. J. Kim: Weierstrass semigroups on a smooth plane curve of degree 4,5 and 6, preprint.
- [KK] S. J. Kim, J. Komeda: Weierstrass points on a non-singular plane curve of degree 7, Research Reports of Kanagawa Institute of Technology **B-31** (2007), 29-34.
- [KM] C. Keem, G. Martens: Curves without plane model of small degree, to appear.
- [KO] C. Keem, A. Ohbuchi: On the Castelnuovo-Severi inequality for a double covering, preprint.

TAKESHI HARUI: DEPARTMENT OF MATHEMATICS, GRADUATE SCHOOL OF SCIENCE, OSAKA UNIVERSITY, TOYONAKA, OSAKA, 560-0043, JAPAN.

E-mail address: takeshi@cwo.zaq.ne.jp, takeshi@gaia.math.wani.osaka-u.ac.jp

KOMEDA JIRYO: DEPARTMENT OF MATHEMATICS, CENTER FOR BASIC EDUCATION AND INTEGRATED LEARNING, KANAGAWA INSTITUTE OF TECHNOLOGY, ATSUGI, KANAGAWA, 243-0292, JAPAN.

E-mail address: komeda@gen.kanagawa-it.ac.jp

AKIRA OHBUCHI: FACULTY OF INTEGRATED ARTS AND SCIENCES, TOKUSHIMA UNIVERSITY, TOKUSHIMA, 770-0814, JAPAN. PARTIALLY SUPPORTED BY GRANT-IN-AID FOR SCIENTIFIC RESEARCH, MINISTRY OF EDUCATION, #17540030

E-mail address: ohbuchi@ias.tokushima-u.ac.jp